

Neutrino Oscillation and Magnetic Moment from $\nu - e^-$ Elastic Scattering.

J. Segura

Max-Planck-Institut für Physik. Werner-Heisenberg Institut. Munich.

and

*Departamento de Ingeniería de Sistemas y Comunicaciones. Escuela Politécnica
Superior. Universidad de Alicante. E-03080 Alicante. Spain*¹

Abstract

We discuss how the measurement of the $\bar{\nu}_e - e^-$ elastic cross section at reactor energies can be used to extract new information on the neutrino oscillation parameters. We also consider the magnetic moment contribution and show how both effects tend to cancel each other when the total cross section is measured; to achieve the separation of each of the effects, experiments capable of measuring angular and energy distributions with respect to the outgoing electron become necessary. We study how their different energy and angle dependence enables such a separation; then, the sensitivity to magnetic moments, masses and mixings is discussed. We also show how these experiments can be sensitive to the magnetic moment of τ neutrinos via $\bar{\nu}_e \leftrightarrow \bar{\nu}_\mu$ oscillation.

Keywords: Neutrino-electron scattering, Neutrino Oscillations, Neutrino magnetic moment, Reactor neutrinos.

PACs numbers: 13.10.+q;13.15.+g;14.60.Pq;13.40.Em.

¹Postal Address. e-mail: segura@evalvx.ific.uv.es, jsegura@disc.ua.es

1 Introduction.

Neutrino-electron elastic scattering is the most elementary purely weak leptonic scattering process; it involves very well known particles (e^-) and neutrinos, whose properties are the subject of a continuous theoretical and experimental study[1]. Thus, this process appears as a crucial probe for the study of neutrino properties.

Nevertheless, the smallness of the cross section seems a serious drawback to obtain precision measurements from this process, specially when low energy neutrinos from reactors are considered; from accelerators, where higher statistics have been achieved, there exist measurements of the weak couplings reaching a 5% precision in the determination of $\sin^2\theta_W$ [2]. In reactors experiments, due to the lower statistics (less events are detected), this achievement has been far from reach until now.

Neutrino-electron elastic scattering experiments have proven to be useful to establish bounds on neutrino electromagnetic properties [3, 4, 5], and, in particular, on neutrino magnetic moments; since the relative contribution of the magnetic moment interaction increases with decreasing values of transferred momentum Q^2 , electrons are the the best target for such a purpose compared to any other, more massive, charged particle. The measurement of the total cross section in $\bar{\nu}_e - e^-$ elastic scattering at reactor experiments give the most restrictive bounds on the magnetic moment of neutrinos (μ_ν) since low energy neutrinos are available and low recoil electrons can be detected ($\mu_{\bar{\nu}_e} < 2.4 \cdot 10^{-10} \mu_B$ from $\bar{\nu}_e e^- \rightarrow \bar{\nu}_e e^-$ [3]). New reactor experiments will be able to improve such bound; for instance, the MUNU experiment will be sensitive to magnetic moments as low as $2 - 3 \times 10^{-11} \mu_B$. In fact, the statistics will be high enough and the control of the background good enough to determine $\sin^2\theta_W$ at the level of a 5%[6]. Apart from the higher statistics, the most outstanding feature of the MUNU experiment is the capability of measuring at some extent both the energy and recoil angle of the electron thus being sensitive to the incoming neutrino energy. Then, this new generation of detectors (MUNU, HELLAZ [7]) will be able to measure at some extent distributions with respect to the energy T and angle θ of recoil electrons.

Such experimental improvement allows the study of other properties of neutrinos, apart from neutrino magnetic moment. As it was shown in ref. [8] the fact that the experiment is able to measure both the angle and energy of the recoil electron, together with the cancellation of the weak cross section $d\sigma^{\bar{\nu}_e}/dT$ (T is the kinetic energy of the recoil electron) for a neutrino energy $E_\nu = m_e/4\sin^2\theta_W$ and forward electrons gives rise to an appearance-like experiment, where the study of neutrino oscillations is available by measuring events around the *dynamical zero* [9]. This novel kind of oscillation experiment was shown to be potentially sensitive to values of masses and mixings not excluded by experiment yet.

In this work, we will estimate the sensitivity of present reactor experiments, taking MUNU as our reference, both for μ_ν and neutrino oscillations. We will show how a disappearance regime for the oscillations is also available and that it

is potentially sensitive to yet unbounded values of the parameters; the oscillation contribution, however, tends to cancel the μ_ν contribution. Then, one should combine data from both regimes to obtain independent bounds on each of the effects; angular (and energy) distribution measurements will be necessary since one can not trust bounds on magnetic moment from total cross sections measurements at least one disregards the possibility of neutrino oscillation. Besides, we will also show how, when $\bar{\nu}_e \leftrightarrow \bar{\nu}_\tau$ oscillation is considered, the experiment is potentially sensitive to combinations of ν_τ magnetic moment and mixings not excluded by experiments so far.

2 Appearance and disappearance regimes.

The cross section when $\bar{\nu}_e \leftrightarrow \bar{\nu}_X$ ($X = \mu, \tau$) oscillation takes place within the distance x from the reactor to the detector reads [8]

$$\frac{d\sigma}{dT}(x, E_\nu, T) = \frac{d\sigma^{\bar{\nu}_e}}{dT} + \sum_i P_{\bar{\nu}_e \rightarrow \bar{\nu}_i}(x) \left(\frac{d\sigma^{\bar{\nu}_\mu}}{dT} - \frac{d\sigma^{\bar{\nu}_e}}{dT} \right) \quad (1)$$

where we have used that $\sum_i P_{\bar{\nu}_e \rightarrow \bar{\nu}_i} = 1$ (we don't consider oscillation to sterile neutrinos) and $d\sigma_{\bar{\nu}_\mu} = d\sigma_{\bar{\nu}_\tau}$ in the Standard Model.

To simplify the analysis, let us consider mixing of two generations ($\bar{\nu}_e \leftrightarrow \bar{\nu}_\mu$), then we should replace:

$$\sum_i P_{\bar{\nu}_e \rightarrow \bar{\nu}_i}(x) \Rightarrow P_{\bar{\nu}_e \rightarrow \bar{\nu}_\mu}(x) = \sin^2(2\phi) \sin^2\left(\frac{\Delta m^2 x}{4E_\nu}\right) \quad (2)$$

where, as usual, Δm^2 is the difference of the squared masses and ϕ the mixing angle.

Taking $E_\nu \simeq m_e/(4\sin^2\theta_W)$ and $\theta \simeq 0$ (θ is the electron scattering angle in the LAB frame) we would have $d\sigma^{\bar{\nu}_e} \ll d\sigma^{\bar{\nu}_\mu}$; then

$$\frac{d\sigma}{dT}(x, E_\nu, T) \simeq \frac{d\sigma^{\bar{\nu}_e}}{dT} + P_{\bar{\nu}_e \rightarrow \bar{\nu}_\mu}(x) \left(\frac{d\sigma^{\bar{\nu}_\mu}}{dT} - \frac{d\sigma^{\bar{\nu}_e}}{dT} \right) > \frac{d\sigma^{\bar{\nu}_e}}{dT} \quad (3)$$

and we see how the kinematics have been chosen to perform an appearance-like experiment in the sense that an excess of events would be detected with respect to the standard model prediction.

This is not the first time $\bar{\nu}_e - e^-$ has been considered in order to study oscillations; nevertheless, ref. [8] is the first one to take advantage of the dynamical zero to perform an appearance-like experiment. The previous analysis [10] were based on the measurement of the total cross section, in a given interval of recoil energies T , integrating over the whole angular range θ (not measured); considering this strategy at reactor energies, the $\bar{\nu}_e$ contribution to (1) after the integration over T and all θ becomes around 2-3 times greater than that from $\bar{\nu}_\mu$, due to

the fact that for $\bar{\nu}_e$ we have both charged and neutral currents and only neutral currents for $\bar{\nu}_\mu$ (CC-NC interference is however negative, as shown explicitly by the appearance of the *dynamical zero*). Then, far from the dynamical zero where $d\sigma^{\bar{\nu}_e} > d\sigma^{\bar{\nu}_\mu}$ (which covers most of the phase space) eq. (1) leads to

$$\frac{d\sigma}{dT}(x, E_\nu, T) \lesssim (1 - P_{\bar{\nu}_e \rightarrow \bar{\nu}_\mu}(x)) \frac{d\sigma^{\bar{\nu}_e}}{dT} < \frac{d\sigma^{\bar{\nu}_e}}{dT} \quad (4)$$

and hence the effect of oscillations would be to reduce the number of detected events when one integrates over T and all θ ; in this case the kinematics have been chosen to perform a disappearance-like experiment.

In this type of disappearance-like experiment, contrary to the appearance case, one can not force one of the two cross section (for $\bar{\nu}_\mu$ or $\bar{\nu}_e$) to be much smaller than the other one; then the relative contribution of the oscillation term will be smaller. But, on the other hand, we have the advantage of higher statistics since integrating for all θ . We will make use of both appearance and disappearance-like regimes to extract information on the oscillation parameters.

3 Estimative bounds on μ_ν and the oscillation parameters.

Let us consider the observable:

$$R(\theta) = \frac{N_o(\theta)}{N_W^e(\theta)} \quad (5)$$

where N_o is the number of events for electron angles lower than θ that would be detected by the experiment considering that oscillations occur and N_W^e is the corresponding standard model prediction; when θ is small and the T -window is located around $T \simeq 2m_e/3$ then $R > 1$ (appearance) while $R < 1$ when we integrate $\forall \theta$ (disappearance).

We will now estimate the bounds one could extract by measuring R , considering for the moment that the statistical error alone accounts for the precision of the experiment, which measures $N_o \pm \sqrt{N_o}$ events. Then, a would-be exclusion plot (1σ) could be derived by setting

$$|R - 1| \equiv \left| \frac{\int P(d\sigma^{\bar{\nu}_\mu} - d\sigma^{\bar{\nu}_e})}{N_W^e} \right| < \frac{\sqrt{N_o}}{N_W^e} \equiv F \quad (6)$$

where F accounts for the precision in the measurement of R .

To make an estimative comparison between both regimes, let us integrate events in the window $0.1 < T < 2 \text{ MeV}$ for two different selection of angles:

a) $\theta < 0.3 \text{ rad}$

b) $\forall \theta$

We will take as N_o (N_W^e, N_W^μ) the total number of events in one year using as normalization the number of events that are expected to be detected by the MUNU experiment in the window $0.5 < T < 2.0 \text{ MeV}$; we take $x = 20 \text{ m}$ (similar to the actual reactor-detector distance in the MUNU experiment).

Using eq. (7) we get the following estimative limits:

	$\Delta m^2 (eV/c)^2 >$ for $\sin^2(2\phi) = 1$	$\Delta m^2 (eV/c)^2 >$ for $\sin^2(2\phi) = 0.35$	$\sin^2(2\phi) <$ for $\Delta m^2 \gg$	Statistical Error
a)	$1.3 \cdot 10^{-2}$	$2.5 \cdot 10^{-2}$	0.21	9.0%
b)	$1.0 \cdot 10^{-2}$	$1.7 \cdot 10^{-2}$	0.04	1.3%

to be compared with those given by disappearance reactor experiments [11]. Note how the disappearance regime is more sensitive to low values of $\sin^2(2\phi)$ ($\langle P \rangle \simeq \frac{1}{2} \sin^2(2\phi)$) than the appearance due to its higher statistics; this fact, contrary to the comparison between "standard" appearance and disappearance experiments, is related the fact that the statistics is higher in the disappearance regime for $\bar{\nu}_e e^-$ elastic scattering. On the other hand, for low Δm^2 we have

$$\langle P \rangle \simeq k(\Delta m^2)^2 \sin^2 2\phi \quad (7)$$

with $k = \langle (x/4E_\nu)^2 \rangle$ and one can check that higher values of E_ν enter in the integration b) compared to integration a); this explains why, being the bound on $\sin^2 2\phi$ for high Δm^2 greater in case a) the bounds for low Δm^2 are similar to those extracted from b). This particular discussion shows the general features of both regimes: similar sensitivity for low Δm^2 but better sensitivity of the disappearance regime to low $\sin^2 2\phi$ (high Δm^2).

The limits extracted from the kinematical selection a) are worse than that for b); but, let us note that to extract all the potentiality of the *dynamical zero*, it will be necessary to consider narrower windows around $T \simeq 2m_e/3$ (and $\theta = 0$) where the cancellation in the standard cross section takes place.

In any case, this first estimation compels us to a closer look to the oscillation effect, since the estimative bounds presented are similar, for large mixing, to those obtained by C.C.-detection while we get lower bounds for the mixing at high Δm^2 for selection b). However, let us remind that, although the statistical error in case b) is quite small we should consider also the systematic error (5%) coming mainly from the lack of precision in the knowledge of neutrino spectrum. Later, we will come back to this point. The sensitivity of the experiment MUNU, following [6], is mainly dominated by this systematic uncertainty and changes very little as a function of the signal versus background ratio.

Let us now estimate which would be the bound on μ_ν that could be extracted from the measurement of events inside the region $0.1 < T < 2.0 \text{ MeV}$, $\forall \theta$. The signature of this contribution would be to increase the total number of expected events since now

$$\frac{d\sigma}{dT} = \frac{d\sigma_e^W}{dT} + \frac{d\sigma_e^M}{dT} > \frac{d\sigma_e^W}{dT} \quad \forall \theta, T \quad (8)$$

Then, considering only the statistical error, if the experiment measures $N \pm \sqrt{N}$ events, being $N = N_W^e + N_{mag}$, one can set a bound on μ_ν ; then, we get $\mu_\nu < 1.3 \cdot 10^{-11} \mu_B$ (1σ) with the integration selection b). If one now integrates only for $\theta < 0.3 \text{ rad}$ (a)) one will have worse statistics (9% compared to 1.3%) but, since the weak cross section becomes smaller, there is a better ratio signal versus weak interaction and one can establish $\mu_\nu < 3.1 \cdot 10^{-11} \mu_B$. Then, at first sight, one could conclude that more stringent bounds, both for the oscillation parameters and for μ_ν are extracted when the total number of events for all angles are measured.

However, comparing eqs. (4) and (8), we see that oscillation effects enter with the opposite sign when measuring events $\forall \theta$ so that both effects could even cancel each other. Then, although the better statistics are achieved integrating over the whole range of energies and angles caution must be taken since the bounds, both for oscillation or neutrino magnetic moment, depend on the assumption of the absence of the other effect. On the other hand, if we now compare eqs. (3) and (8), we see how both effects tend to increase the total number of events. In fact, on the *dynamical zero* ANY extra contribution will be additive, if it appears, since the standard weak cross section is zero; then one could set bounds over one of such non-standard effects neglecting the rest. In particular, for $0.1 < T < 0.5 \text{ MeV}$ and $\theta < 0.3 \text{ rad}$, under the same assumptions as before we get

	$\Delta m^2 (eV/c)^2 \geq$ for $\sin^2(2\phi) = 1$	$\Delta m^2 (eV/c)^2 \geq$ for $\sin^2(2\phi) = 0.35$	$\sin^2(2\phi) \leq$ for $\Delta m^2 \gg$	Statistical Error
a)	$8.1 \cdot 10^{-3}$	$1.5 \cdot 10^{-2}$	0.35	60%
b)	$9.3 \cdot 10^{-3}$	$1.6 \cdot 10^{-2}$	0.05	1.7%

while, in the same region, we get the bound $\mu_\nu < 2.8 \cdot 10^{-11} \mu_B$ for a) and $\mu_\nu < 1.2 \cdot 10^{-11} \mu_B$ for b). With this selection in the T -window we obtain, with a much higher statistical error, bounds on oscillations parameters somewhat better than those from disappearance-like for high of $\sin^2 2\phi$ and similar bounds for μ_ν but worse bounds for $\sin^2 2\phi$ at high Δm^2 . In any case, it seems that both appearance and disappearance regimes can potentially give relevant information on the oscillation effect and that similar bounds on μ_ν can be given with both kinds of integration region; unfortunately, as commented before, the disappearance region shows a clear disadvantage: μ_ν and oscillation effects tend to cancel; this fact is

illustrated in figure 1 where the σ -deviation of the observable R, both for oscillation and magnetic moment, with respect to 1 is plotted as a function of the angle of integration; in this figure the parameters are fixed to one of the bounds extracted from the integration $\forall \theta$ (b)).

Then, as a summary of our previous estimations, and to fix ideas, let us underline some facts:

- 1.- Both regimes are sensitive to $\bar{\nu}_e \leftrightarrow \bar{\nu}_x$ where $\bar{\nu}_x$ stands for any non-sterile neutrino; there is no production threshold for the detection.
- 2.- Similar bounds for large $\sin^2 2\phi$ are extracted from appearance and disappearance regimes, but disappearance gives better bounds for small mixing (neglecting μ_ν interaction). The statistical error is much larger in the appearance regime.
- 2.- Disregarding oscillations, μ_ν is better measured integrating $\forall \theta$.
- 3.- Disappearance-like contributions could mask μ_ν interaction when an integration $\forall \theta$ is performed.
- 4.- Any non-standard effect would increase the number of detected events around the dynamical zero ($T \simeq 2m_e/3$ and $\theta = 0$).

4 Disentangling the magnetic moment effect from oscillation.

Only if an experiment is able to measure both the recoil angle of the electron θ and its energy T one can safely separate both effects, since, as explained, though they tend to cancel each other when total cross sections are measured, the angular (and energy) distributions for oscillation and μ_ν -interaction are different (the μ_ν term is always additive and oscillations add near the *dynamical zero* but they subtract far away from it); furthermore, one can conclude from the previous estimations that both regions (far away and close to the *dynamical zero*) are sensitive to oscillations and magnetic moment. The one could fit the shape of the distributions to separate both effects; a deviation from the expected θ dependence would be mainly a signal of oscillation while μ_ν would be observed as an overall excess of events. One can also construct convenient observables to achieve such a separation; for instance, and as an illustration, let us consider the quantity:

$$\mathcal{O}_o = \frac{N_o(\theta < \theta_0)}{N_o(\theta > \theta_0)} \bigg/ \frac{N_W(\theta < \theta_0)}{N_W(\theta > \theta_0)} \quad (9)$$

where, given a window in T , $N(\theta < \theta_0)$ is the number of observed events N_o and of expected events from the S.M. predictions (N_W) that would be detected for

angles lower than θ_0 ; $N(\theta > \theta_0)$ is the corresponding number of events for angles higher than θ_0 .

Considering an observable such like $\mathcal{O}_o$ will have several benefits: first, we can take advantage of both the apparition ($N(\theta < \theta_0)$) and disappearance regimes ($N(\theta > \theta_0)$) for the oscillation contribution; second, since we are integrating in the same window of T both for $\theta < \theta_0$ and $\theta > \theta_0$ similar neutrino energies appear, thus canceling neutrino spectrum uncertainties partially and total flux uncertainty completely; and third, due to their different angular dependence the observable will be able to separate oscillation contributions from magnetic moment contributions.

With all this arguments in mind, let us check what is the σ deviation of $\mathcal{O}$ from 1 (value without non-standard effects) for different values of θ_0 and different selections in the window in T ; since we expect some dependence $\mathcal{O}(\theta_0)$ we will also estimate the error in the measurement of θ , taking $\mathcal{E}(\theta) = 0.05 \text{ rad}$ [6]; we will consider that the precision is given mainly by the statistical error and by the uncertainty in the determination of θ and thus in the actual value of the observable $\mathcal{O}_o(\theta_0)$ (systematic error); we sum in quadrature both uncertainties. Fig. (2) shows the sigma deviations of $\mathcal{O}_{osc}$ for different selection of T -windows and oscillation parameters. In the same figure, we also plot the σ -deviation of the observable for electromagnetic interaction taking $\mu_\nu = 2.3 \cdot 10^{-11} \mu_B$

From fig. (2) we observe that the region of low $\theta \sim 0.3 \text{ rad}$ is mainly sensitive to oscillation; in this region the observable $\mathcal{O}$ is not much sensitive to μ_ν ; also, let us notice that similar bounds for high $\sin^2 2\phi$ to those from the measurement of R can be extracted (both appearance and disappearance were equally sensitive to this values) while the bounds on $\sin^2 2\phi$ at high Δm^2 are between those extracted from the appearance and disappearance regimes, improving the first ones; for instance, in the window $0.1 < T < 0.5 \text{ MeV}$ and for $\sin^2(2\phi) = 1$ one gets $\Delta m^2 < 10^{-2} \text{ eV}^2$ at $\theta_0 = 0.3 \text{ rad}$ with a 40% error in the measurement of $\mathcal{O}$ (curve 1, dashed) while for the window $0.1 < T < 2.0 \text{ MeV}$ and large Δm^2 one can set $\sin^2(2\phi) < 0.15$ with a 10% error (curve 2, solid).

On the other hand, for large $\theta \sim 1 \text{ rad}$ we have a better sensitivity to μ_ν , which also lies between the bounds extracted from the selections a) and b) for the angular region ($\mu_\nu < 2.3 \cdot 10^{-10} \mu_B$ for $0.1 < T < 2.0 \text{ MeV}$ with a 6% error. Then, we see that there are two regions and each of them is mainly sensitive to one of the effects considered. Besides, for small angles, the mag. moment effect is of the same sign as the oscillation one so that there is no risk of cancellation between them. Notice also the dependence of the effects on the selection of T -window; mag. moment. contribution and oscillation for large Δm^2 are better measured when the T -window is large (since in this case the most important thing is to get as high statistics as possible, as it was shown previously) while for large $\sin^2 2\phi$ and small Δm^2 it is more convenient to chose a narrower window around the dynamical zero (in our case $0.1 < T < 0.5 \text{ MeV}$). Then, this is a tunable experiment which can prospect different regions of the parameter space

by choosing different values of the angle and recoil energy.

5 $\bar{\nu}_\tau$ electromagnetic properties from $\bar{\nu}_e - e^-$ elastic scattering.

Finally, it is interesting to note that, in case oscillations take place, any extra interaction of the new flavors originated from oscillations, could affect the value of the elastic cross section; consider, for instance, that $\bar{\nu}_e$ oscillate to $\bar{\nu}_\tau$ and that the tau neutrino has a large magnetic moment (the best lab. bound is $4 \cdot 10^{-6} \mu_B$ [12]). To simplify the analysis, let us consider that the magnetic moment cross section for the tau neutrino is much larger than the weak cross sections (which is granted for $\mu_\nu > 10^{-9} \mu_B$) and that τ -neutrino mass effects are negligible in the cross section; then, we would expect an excess to respect to the standard model prediction and one can write

$$\frac{d\sigma}{dT} \simeq \frac{d\sigma^{\bar{\nu}_e}}{dT} + \mu^2 P(x) \frac{\pi \alpha^2}{m_e^2} \frac{1 - T/E_\nu}{T} ; \mu = \mu_{\bar{\nu}_\tau} / \mu_B \quad (10)$$

where the terms $P(x)d\sigma^{\bar{\nu}_e}$ and $P(x)d\sigma^{\bar{\nu}_\tau}$ have been neglected. Then one gets the following 1σ bounds from the observable $\mathcal{O}$:

- $\chi^2 \sin^2(2\phi) < 0.1$ at large Δm^2 for $0.1 < T < 2.0 \text{ MeV}$, $\theta \sim 1 \text{ rad}$; total error $\sim 5\%$.
- $\chi^2 \sin^2(2\phi) (\Delta m^2)^2 < 2 \cdot 10^{-5} (eV^4)$ at large $\sin^2(2\phi)$ for $0.1 < T < 0.5 \text{ MeV}$, $\theta \sim 0.4 \text{ rad}$; total error $\sim 20\%$.

where $\chi = \mu_{\bar{\nu}_\tau} / 10^{-10} \mu_B$.

This kind of bounds, although relating unknown parameters, would explore combinations of values which are not excluded yet; or, in other words, the elastic neutrino-electron cross section is sensitive to non standard neutrino physics in still admissible scenarios.

6 Conclusions.

We should stress again that the measurement of the total cross section in present reactor experiments is potentially sensitive both to values of μ_ν and the oscillation parameters not excluded yet. By measuring the total cross section one can not set bounds on μ_ν neglecting oscillation since both effects tend to cancel. Experiments able to measure energy and angle of recoil electrons are needed; only then one can safely separate both effects and set independent bounds on each of them. Furthermore, the elastic neutrino-electron cross section is sensitive to

non-standard neutrino interactions induced by oscillation, as is the case of magnetic moment interaction of $\bar{\nu}_\tau$ ($\bar{\nu}_\mu$), for still available values of the parameters. Therefore, this process promises a better understanding of neutrino dynamics.

Acknowledgements.

The author wishes to acknowledge the support from *Fundación Ramón Areces* for his postdoctoral fellowship. I also wish to thank G. Raffelt for his hospitality during my stay at M.P.I. für Physik-Munich and M.C. González-García, L.M. García-Raffi and A. Gil for useful discussions.

References

- [1] K. Winter, Lepton-Photon Symp.(1995)569.
G. G. Raffelt, Blois 1992, Proceedings, Particle Astrophysics 99-110.
- [2] P. Vilain et al., Phys. Lett B335(1994)246.
- [3] G.S. Vidyakin et al., JETP Lett. 49(1989)740; JETP Lett. 55 (1992) 206.
- [4] D.A. Krakauer et al., Phys. Lett. B252(1990) 177;
P. Vilain et al., Phys. Lett. B345(1995)115.
- [5] J. Bernabéu, S.M. Bilenky, F.J. Botella y J. Segura, Nucl. Phys. B246(1994)434.
- [6] C. Brogini et al., LNGS-92/47 (MUNU proposal).
- [7] F. Arzarello et al., Preprint CERN-LAA-94-19.
- [8] J. Segura, J. Bernabéu, F.J. Botella y J.A. Peñarrocha, Phys. Lett. B335(1994)93.
- [9] J. Segura, J. Bernabéu, F.J. Botella y J. Peñarrocha, Phys. Rev. D49(1994)1633.
- [10] W.J. Marciano, Phys. Rev. D36(1987)2859;
S.P. Rosen, B. Kayser, Phys. Rev. D23(1981)669;
L.A. Ahrens et al., Phys. Rev. D31(1985)2732.
B. Halls, B.H.J. McKellar, Phys. Rev. D24(1981) 1785.
- [11] B. Achkar et al., Nucl. Phys. B434(1995)503;
G. Zacek et al., Phys. Rev. D34(1986)2621;
G.S. Vidyakin et al., JETP Lett. 59(1984)364.
- [12] H. Grotch, R. Robinet, Z. Phys. C39(1988) 563;
T.M. Gould, I.Z. Rothstein, Phys. Lett. B333(1994)545.

Figure captions:

Figure 1: σ -deviation of the observable $R(\theta)$ from 1 for $0.1 < T < 0.5 \text{ MeV}$ (dashed) and $0.1 < T < 2.0$ (solid). $\Delta m^2 = 10^{-2} eV^2$ and $\sin^2 2\phi = 1$ for curves **1**; $\Delta m^2 = 1 eV^2$ and $\sin^2 2\phi = 0.04$ for **2**; $\mu_\nu = 1.2 \cdot 10^{-11} \mu_B$ for **3**.

Figure 2: Sigma-deviation of the observable $\mathcal{O}_o(\theta_0)$ from 1 for $0.1 < T < 0.5 \text{ MeV}$ (dashed) and $0.1 < T < 2.0 \text{ MeV}$ (solid). $\Delta m^2 = 10^{-2} eV^2$ and $\sin^2 2\phi = 1$ for curves **1**; $\Delta m^2 = 1 eV^2$ and $\sin^2 2\phi = 0.15$ for **2**; $\mu_\nu = 2.3 \cdot 10^{-11} \mu_B$ for **3**.




Figure 1




Figure 2